

Leishmaniasis in Bolivia. I. *Lutzomyia longipalpis* (Lutz & Neiva, 1912) as the vector of visceral leishmaniasis in Los Yungas

F. LE PONT¹ AND P. DESJEUX²

¹ORSTOM, IBBA, c/o Embajada de Francia, Casilla 824, La Paz, Bolivia; ²Institut Pasteur Paris, IBBA, c/o Embajada de Francia, casilla 824, La Paz, Bolivia

Abstract

A relatively high leishmanial infection rate was found in the phlebotomine sandfly *Lutzomyia longipalpis* collected from three villages of the Los Yungas region (Department of La Paz, Bolivia). 2,578 female sandflies were dissected. In three houses surveyed in Santa Barbara promastigote infection rates of *Lu. longipalpis* were 4.2, 2.2 and 3.2% respectively. Anatomical localization of the infection in the insect, and biochemical characterization of the strains indicate that the parasite belongs to the *Leishmania donovani* complex. The geographical area and the biotopes of *Lu. longipalpis* are discussed in relation to the vector-parasite relationship.

Introduction

Epidemiological studies on kala-azar in neo-tropical regions have been carried out mainly in the north-east and north of Brazil, where the sandfly *Lutzomyia longipalpis* is considered to be the vector of the disease (DEANE, 1956; DEANE & DEANE, 1964). However, since the pioneer works of CHAGAS *et al.* (1937, 1938), the presumption that this sandfly plays a leading role in the transmission of kala-azar has remained based only on indirect evidence (DEANE & DEANE, 1954b; LAINSON *et al.*, 1977a, LAINSON & SHAW, 1979; LAINSON *et al.*, 1983). This evidence is: (i) the predominance of *Lu. longipalpis* in most foci of human and canine visceral leishmaniasis, (ii) the same distribution of the insect and the human disease throughout Latin America, (iii) the distribution around human habitations and endophily, (iv) the attraction to man, (v) the experimental infection obtained from infected patients, dogs and foxes, (vi) the experimental transmission of the parasite from hamster to hamster by the bite of infected *Lu. longipalpis*, and (vii) the elimination of the disease after systematic spraying of houses with insecticides.

Only a few studies of the natural infection rate in *Lu. longipalpis* have been made to date (DEANE & DEANE, 1954a; RYAN *et al.*, 1984; LAINSON *et al.*, 1985) and biochemical characterization of the strains has not been completed but it seems that the *L. donovani* complex may be implicated in spontaneous infections.

DEANE & DEANE (1954a) studied a focus of visceral leishmaniasis in the suburbs of Sobral (Ceara), Brazil, situated at an altitude varying from 100 to 800 metres.

At the end of December, just before the rainy season, they found two of 141 *Lu. longipalpis* dissected to be infected (1.4%). The infection was heavy with a supra-pylarian localization, similar to infection by *L. chagasi* s.l. (LAINSON *et al.*, 1977b). A third specimen was found among 876 additional dissected *Lu. longipalpis*, giving an infection rate of 0.3% (1017 sandflies examined).

During an epidemic outbreak, LOPES (1956) found in Jacobina (Bahia) eight of 209 (3.8%) *Lu. longipalpis* infected with promastigotes: four of the infected specimens came from chicken coops, three from pig-sties and one from a tree trunk sheltering hens. In

the same city, SHERLOCK & PESSOA (1966) and SHERLOCK & GUITTON (1969) found one *Lu. longipalpis* infected of 91 samples examined.

RYAN *et al.* (1984) have recently recorded the presence of heavy suprapylarian promastigote infections in 8 of 1,500 (0.5%) *Lu. longipalpis* on the island of Marajó, Pará.

In 1984, LAINSON *et al.* (1985) in the focus of Santarém, Pará State, found 35 (7.14%) of 491 *Lu. longipalpis* dissected to be infected, and the parasite was transmitted to hamsters by the bites of infected wild flies.

From October 1982, epidemiological studies on the vector of visceral leishmaniasis in Bolivia have been carried out. In the Los Yungas area (Department of La Paz), both canine and human visceral leishmaniasis were described recently (ANGLES *et al.*, 1982; DESJEUX *et al.*, 1983), confirming a suspicion that kala-azar may occur in this region (GATTI *et al.*, 1939). VELASCO (1973) surveyed the phlebotomine sandfly fauna during the cool and dry winter season (from June to August); he noted, for the first time, the presence of *Lu. longipalpis* and found it very common and widespread (present in 65% of collections).

These observations led us to investigate the rate of natural infection among the population of *Lu. longipalpis* in this area.

Materials and Methods

Area studied

Our area of investigation (see Fig.) is located in the village of Santa Barbara, near the village of Chijchipa, where a second case of kala-azar was described in November 1982 (Urjel & Desjeux, 1982, personal communication). Santa Barbara is situated at the foot of a valley overlooked to the south-east by the village of Coroico, the county-town of Nor-Yungas Province.

The Yungas constitute a very peculiar geographical terrain. They are steep-sided valleys in the foothills of the Eastern Andean Cordillera. Santa Barbara, at an altitude of 950 m, is overlooked by the mountains of Uchumachi and Kusilluni, rising to a height of 2500 metres. The valley of Coroico River is directed north-east. The climate is subtropical with an average yearly temperature of 18°C, and a mean annual rainfall of 1200mm. Some of the inhabitants are black and live under poor conditions, cultivating coffee, coca plants and citrus trees. Steep slopes going down from Coroico are covered with wet subtropical forest. Coffee




Fig. Map showing the area surveyed.

plantations and cultivated plots surround the villages and are situated at the bottom of the valleys. This area is less populated and much more humid than that of Coripata/Khala-Khala, where canine visceral leishmaniasis was described (ANGLES *et al.*, 1982).

Three villages were surveyed: Santa Barbara (950 m), with houses scattered all along the Coroico River; Tocania (1320 m), on the north-western slope near Chijchipa; and Cedro Mayo (1370 m), situated on a slope, 9 km from Santa Barbara.

Sandfly collections

Phlebotomine sandflies were collected in October and November 1982, at the beginning of the rainy season. Collections, using human bait, were made between 19 and 24 hours, around isolated houses.

Isolation of *Leishmania* from sandflies

For all sandflies collected, species were identified by the examination of genitalia. Females were dissected; when found infected by promastigotes, the intestinal tract was picked up from the slide and carefully disrupted in 0.5 ml of sterile 9% NaCl. The suspension was thrown into modified NNN rabbit blood agar base (DECKER-JACKSON & HONBERG, 1978), with a (40% v/v, overlay/base) liquid overlay of PBS.

The tubes were examined every week and discarded if negative after one month. The isolations were routinely passaged.

Isoenzyme characterization of the stocks

Multiplication of the stocks was obtained dividing the contents of a single tube between larger tubes; cultures were routinely harvested, centrifuged at 2500 rpm for 10 min and washed (three times); pellets were stored at -80°C until use. Just before electrophoresis, an equal volume of hypotonic enzyme stabilizer was added. Lysis of the parasites was achieved through freeze-thawing procedures repeated three times.

The following enzymic systems were used for characterization:

Malate dehydrogenase E.C.1.1.1.37 (MDH)

Malate dehydrogenase (oxaloacetate decarboxylating) NADP⁺ E.C.1.1.1.40 (ME)

Isoctrate dehydrogenase E.C.1.1.1.42 (ICD)

6 Phosphogluconate dehydrogenase E.C.1.1.1.44 (6PGDH)

Glucose 6 phosphate dehydrogenase E.C.1.1.1.49 (G6PDH)

Phosphoglucomutase E.C.2.7.5.1. (PGM)

Mannose phosphate isomerase E.C.5.3.1.8. (MPI)

Glucose phosphate isomerase E.C.5.3.1.9. (GPI)
Glutamate dehydrogenase E.C.1.4.1.2. (GDH/NAD and GDH/NADP⁺)
Aconitate hydrolase E.C.4.2.1.3. (ACON)
Peptidase E.C.3.4.11.11. (PEP)

Five reference strains were used:

ITMAP 263 *L. infantum* (man)—Morocco, 1967.

M 2682 *L. chagasi* s.l. (man)—Bahia State, Brazil, 1974

M 2904 *L. braziliensis braziliensis* (man)—Pará State, Brazil, 1975.

PH 8 *L. mexicana amazonensis* (*Lu. flaviscutellata*)—Pará State, Brazil, 1967

M 5088 *L. hertigi deanei* (*Coendou prehensilis*)—Pará State, Brazil, 1978.

The electrophoresis were carried out on cellulose acetate plates (Helena Laboratories) running at 200 volts for 30 min. All cell buffers were used with 20% sucrose, their preparation being adapted from LANHAM *et al.* (1981) and as previously described (TIBAYRENC & DESJEUX, 1983).

Results

Infection rate of *Lu. longipalpis*

Lu. longipalpis (see Table) represented 96% of all the phlebotomine sandflies captured. 2578 females were dissected. Promastigote infection rates of *Lu. longipalpis* were, 4.2, 2.2 and 3.2% respectively in three houses surveyed in Santa Barbara; in the fourth, a nearby poultry farm with walls covered with a coating containing residual insecticide, no infected sandflies were found.

Two *Lu. longipalpis* of 240 (0.8%) were found infected in a house located 300 metres below the village of Tocania. Very similar results were obtained in Cedro Mayo: two specimens out of 187 were infected (1%).

In every infected sandfly, the parasitism was heavy, with distension of the upper digestive track (proventriculus and cardia). The localization of the promastigotes in the suprapyloric region together with their large size, suggest that the parasite belongs to the *L. donovani* complex (LAINSON *et al.*, 1977b).

Isolation of *Leishmania* from *Lu. longipalpis* specimens

Five strains were isolated from infected *Lu. longipalpis* females, and then subcultured weekly. Semi-

Table I—Results of catches and dissections of *L. longipalpis* achieved around human habitations in the focus of visceral leishmaniasis in Los Yungas (Coroico village), Department of La Paz. October/December 1982. Dwellings (1) (3) (5) and (6) had dogs.

Village	Santa Barbara				Tocania	Cedro Mayo
Altitude	950 m				1320 m	1370 m
Situation	Bottom of valley				North-west slope of valley	As Tocania but 9 km upstream
	(1) Dom Angel family	(2) Adriana family	(3) Nicolas family	(4) Poultry farm	(5) Sabala family	(6) Pinedo family
No. of <i>L. longipalpis</i> dissected	1910	45	91	105	240	187
No. of <i>L. longipalpis</i> infected (+)	81	1	3	—	2	2
Rate of infection	4.2%	2.2%	3.2%	—	0.8%	1.0%

massive cultures were obtained, giving sufficient material for enzyme electrophoresis.

Isoenzyme characterization of the stocks

The electrophoretic result for the five Bolivian stocks studied showed, for 12 enzymes, a consistent enzymic profile, similar to that of ITMAP-263 (*L. infantum*) and M-2682 (*L. chagasi* s.l.); they were easily distinguished from *L. m. amazonensis*, *L. b. braziliensis* and *L. h. deanei*. We conclude that the five strains of *Leishmania* isolated from *Lu. longipalpis* specimens belong to the *L. donovani* complex.

Further details of these enzyme similarities between *Lu. longipalpis* Bolivian stocks, one stock from a human Bolivian visceral case, and visceral *Leishmania* reference stocks from Brazil and Morocco, will be published elsewhere (DESJEUX *et al.*, in press), discussing their significance and taxonomic consequences.

Discussion

The Yungas region of Bolivia is well known as a focus of mucocutaneous leishmaniasis or espundia (DESJEUX, 1974, 1976; DESJEUX *et al.*, 1974), which affects a large proportion of the inhabitants. The discovery in 1982 of two cases of visceral leishmaniasis in the same area, associated with widespread canine visceral leishmaniasis, together with the finding of numerous specimens of *Lu. longipalpis* with a relatively high infection rate, provokes evidence that in this region transmission cycles of both visceral and mucocutaneous leishmaniasis co-exist.

The geography is favourable and this area, with its mild climate but with an Amazonian influence, presents an extraordinary diversity of biotopes. However, the visceral leishmaniasis focus studied shows the general characteristics observed in other neotropical foci, i.e., (i) villages with scattered houses, (ii) poor agriculture on small plots of land, with secondary vegetation, (iii) part of the population being black (negro), as in the foci of Venezuela (PIFANO *et al.*, 1976) or Brazil (VIANNA MARTINS *et al.*, 1956; SHERLOCK & PESSOA, 1966); (iv) location in warm foothills in which DEANE & DEANE (1964) had noted (as *bosqueirao* or *pe da serra*) that *Lu. longipalpis* infection rates are higher in the valleys than on the slopes; and (v) in our three foci, peri-domestic phlebotomine sandflies represent a considerable inconvenience to the inhabitants, with *Lu. longipalpis* predominant. We have observed that the populations of this species are particularly aggressive in the spring (September to December), as observed by ARJONA (1971) in Colombia.

However, in spite of the fact that most features in our foci are similar to others in South America, two peculiarities may be noted in this focus—the altitude level of 1100-1600 metres, and the co-existence of both visceral and mucocutaneous leishmaniasis, a fact rarely mentioned in the literature so far.

Acknowledgements

The authors wish to thank Drs. R. Lainson and J. J. Shaw, from the Instituto Evandro Chagas da Fundação SESP, Belém, Pará, Brazil, and Prof. Dr. D. Le Ray, from the Institut de Médecine Tropicale "Prince Léopold", Anvers, Belgium, for providing the *Leishmania* reference stocks.

This research was supported by grants from the French Ministry of Co-operation (STD Service), and a grant from

the UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases.

References

- Angles, R., Le Pont, F. & Desjeux, P. (1982). Visceral canine leishmaniasis in Bolivia. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **76**, 704.
- Arjona, A. C., Osorno-Mesa, E. & Marquez, J. P. (1971). Contribución al estudio epidemiológico del kala-azar en Colombia. *Revista de la Facultad de Medicina, Universidad Nacional de Colombia*, **37**, 90-94.
- Chagas, E., Cunha, A. M., Castro, G. O., Ferreira, L. C. & Romaña, C. (1937). Leishmaniose visceral americana. Relatório dos trabalhos realizados pela Comissão encarregada do estudo sobre a leishmaniose visceral americana em 1936. *Memórias do Instituto Oswaldo Cruz*, **32**, 321-389.
- Chagas, E., Cunha, A. M., Ferreira, L. C., Deane, L., Deane, G., Guimarães, F. N., Paumgarten, M. J. & Sá, B. (1938). Leishmaniose visceral americana. Relatório dos trabalhos realizados pela Comissão encarregada do estudo da leishmaniose visceral americana em 1937. *Memórias do Instituto Oswaldo Cruz*, **33**, 89-229.
- Deane, L. M. (1956). *Leishmaniose visceral no Brasil*. Serviço Nacional de Educação Sanitária, Rio-de-Janeiro, Brasil.
- Deane, L. M. & Deane, M. P. (1964). Leishmaniose visceral nas Américas do Sul e Central. *Arquivos de Higiene e Saúde Pública (São Paulo)*, **29**, 89-94.
- Deane, M. P. & Deane, L. M. (1954). Infecção natural do *Phlebotomus longipalpis* por leptomonas, provavelmente de *Leishmania donovani*, em foco de calazar, no Ceará. *Hospital (Rio-de-Janeiro)*, **45**, 697-702.
- Deane, M. P. & Deane, L. M. (1954b). Infecção experimental do *Phlebotomus longipalpis* em rapôsa (*Lycalopex vetulus*) naturalmente parasitada pela *Leishmania donovani*. *Hospital (Rio-de-Janeiro)*, **46**, 651-653.
- Decker-Jackson, J. E. & Honiberg, B. M. (1978). Glycoproteins released by *Leishmania donovani*: immunologic relationship with host and bacterial antigens and preliminary biochemical analysis. *Journal of Protozoology*, **25**, 514-525.
- Desjeux, P. (1974). *Leishmaniose cutanée et cutanéomuqueuse américaine. Etude de 113 cas observés en Bolivie*. Thèse Doctorat de Médecine, Paris, 132 pp.
- Desjeux, P. (1976). Relations leishmaniose et altitude en Bolivie: Formes cliniques, données épidémiologiques. *Colloque INSERM: "Anthropologie et Biologie des Populations Andines"*, Toulouse 1976, Editions INSERM, pp. 247-256.
- Desjeux, P., Aranda, E., Aliaga, O. & Mollinedo, S. (1983). Human visceral leishmaniasis in Bolivia: first proven autochthonous case from Los Yungas. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **77**, 851.
- Desjeux, P., Quilici, M. & Lapiere, J. (1974). A propos de 113 cas de leishmaniose cutanée et cutanéomuqueuse observés en Bolivie. Etude séro-immunologique de 71 cas. *Bulletin de la Société de Pathologie Exotique*, **67**, 387-395.
- Gatti, G., Boggino, J. & Prieto, C. (1939). Un nouveau foyer de leishmaniose viscérale en Amérique du Sud. *Bulletin de la Société de Pathologie Exotique*, **32**, 602-605.
- Lainson, R. & Shaw, J. J. (1979). The role of animals in the epidemiology of South American leishmaniasis. In: *The Biology of the Kinetoplastidae*, Vol. 2, Lumsden, W. H. R. & Evans, D. A. (Editors). London: Academic Press, pp. 1-116.
- Lainson, R., Ward, R. D. & Shaw, J. J. (1977a). Experimental transmission of *Leishmania chagasi*, causative agent of neotropical visceral leishmaniasis, by the sandfly *Lutzomyia longipalpis*. *Nature*, **266**, 628-630.
- Lainson, R., Ward, R. D. & Shaw, J. J. (1977b). *Leishmania* in phlebotomid sandflies. VI. Importance of hindgut development in distinguishing parasites of the *Leishmania mexicana* and *L. braziliensis* complexes. *Proceedings of the Royal Society (B)*, **199**, 309-320.

- Lainson, R., Shaw, J. J., Silveira, F. T. & Fraiha, H. (1983). Leishmaniasis in Brazil. XIX. Visceral leishmaniasis in the Amazon Region, and the presence of *Lutzomyia longipalpis* on the island of Marajó, Pará State. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **77**, 323-330.
- Lainson, R., Shaw, J. J., Ryan, L., Ribeiro, R. S. M. & Silveira, F. T. (1985). Leishmaniasis in Brazil. XXI. Visceral leishmaniasis in the Amazon region and further observations on the role of *Lutzomyia longipalpis* (Lutz & Neiva, 1912) as the vector. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **78**.
- Lanham, S. M., Grendom, J. M., Miles, M. A., Póvoa, M. M. & Souza, A. A. A. (1981). A comparison of electrophoretic methods of isoenzymes characterization of trypanosomatids. I: Standard stocks of *Trypanosoma cruzi* zymodemes for northeast Brazil. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **75**, 742-750.
- Lopes, J. A. S. (1956). *Phlebotomus longipalpis* naturalmente infectados com formas em leptomonas na cidade de Jacobina, Estado da Bahia. *Revista de Medicina do Parana*, **25**, 57-58.
- Pifano, F. C., Rodriguez, A. A., Romero, J. M., Ortiz, I. & Alvarez, A. (1976). Sobre un nuevo foco de leishmaniasis visceral en el area de Chuao, Estado Aragua, Venezuela. Características bioecológicas y epidemiológicas. *Revista del Instituto Nacional de Higiene*, **9**, 97-105.
- Ryan, L., Silveira, F. T., Lainson, R. & Shaw, J. J. (1984). Leishmanial infections in *Lutzomyia longipalpis* and *Lu. antunesi* (Diptera: Psychodidae) on the island of Marajó, Pará State, Brazil. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **78**, 547-548.
- Sherlock, I. A. & Guitton, N. (1969). Observações sobre calazar em Jacobina, Bahia. III. Alguns datos sobre o *Phlebotomus longipalpis*, o principal transmissor. *Revista Brasileira de Malariologia e Doenças Tropicais*, **21**, 541-548.
- Sherlock, I. A. & Pessôa, S. B. (1966). *Leptomonas* infectando naturalmente *Phlebotomus* em Salvador (Bahia, Brasil). *Revista Latino-Americana de Microbiologia e Parasitologia*, **8**, 47-50.
- Tibayrenc, M. & Desjeux, P. (1983). The presence in Bolivia of two distinct zymodemes of *Trypanosoma cruzi*, circulating sympatrically in a domestic transmission cycle. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **77**, 73-75.
- Vianna-Martins, A., Brener, Z., Mourao, O. G., Moura Lima, M., De Souza, M. A. & Da Silva, J. E. (1956). Calazar autoctone em Minas Gerais. *Revista Brasileira de Malariologia e Doenças Tropicais*, **8**, 555-563.
- Velasco, J. E. (1973). *The phlebotomine sandflies of the Los Yungas region of Bolivia*. M.S. Thesis, Louisiana State University, Department of Tropical Medicine and Parasitology, 204 pp.

Accepted for publication 25th June, 1984.



The First International Seminar on

DENGUE HAEMORRHAGIC FEVER in the Americas

San Juan, Puerto Rico—June 15-16, 1985

Sponsored by the Pan American Health Organization, the Puerto Rican Department of Health and the Centers for Disease Control.

For further information contact: Dr. George Kuno, Centers for Disease Control, Center for Infectious Diseases, San Juan Laboratories, GPO Box 4532, San Juan, Puerto Rico 00936. Telephone: 781-3636; Cables: CDCLABS.