

Ann. Soc. belge Méd. trop.

1985, 65, Suppl. 1, 59-61

ISOZYMIC VARIABILITY OF TRYPANOSOMA CRUZI: BIOLOGICAL
AND EPIDEMIOLOGICAL SIGNIFICANCE

by

M. TIBAYRENC¹, F. BRENIERE¹, C. BARNABE², J.-L. LEMESRE³,
L. ECHALAR⁴ & P. DESJEUX⁵

¹ORSTOM, IBBA, Embajada de Francia, Casilla 824, La Paz, Bolivia

²IBBA, Embajada de Francia, Casilla 824, La Paz, Bolivia

³CIBP, Institut Pasteur, 59019 Lille, France

⁴Universidad Mayor de San Andrés, IBBA, Casilla 841, La Paz, Bolivia

⁵Institut Pasteur de Paris, IBBA, Embajada de Francia, Casilla 824, La Paz, Bolivia

Summary - Genetic interpretation of *T. cruzi* zymograms has led to various hypotheses which are reviewed.

KEYWORDS: Trypanosoma cruzi; Zymograms; Genetic Interpretation.

Isozymic studies on *Trypanosoma cruzi* were initiated by Toyé (1974) and largely developed by Miles et al. (1977, 1980). These authors, by means of a phenetic interpretation of *T. cruzi* zymograms, distinguished in Brazil 3 main isozymic strains («zymodemes) which were classified by numerical taxonomy using isozymic banding (Ready and Miles, 1980) Genetic interpretation of *T. cruzi* zymograms was performed by comparison with better known organisms and led us to the following hypotheses.

Diploidy

T. cruzi zymograms exhibit some typical heterozygous patterns for a diploid organism (Tibayrenc et al., 1981a). By DNA measurement, we verified that the ploidy is constant among all the isozymic strains (Lemesre and Tibayrenc, 1983). This fact was also verified with isozymic patterns (Tibayrenc et al., 1985). Diploid structure was inferred by Tait (1980) for *T. brucei* with similar arguments. Lanar, Levy and Manning (1981), by a DNA study, proposed also the hypothesis of diploidy for *T. cruzi*.

Enzyme quaternary structure

Heterozygous patterns reveal a monomeric structure for phosphoglucomutase, a dimeric structure for isocitrate deshydrogenase, glucose phosphate isomerase, 6-phosphogluconate deshydrogenase, the two loci of malic enzyme, and a pentameric structure for glutamate deshydrogenase Nadp + (Tibayrenc et al., 1981a and 1985).

Lack of Mendelian sexuality at present

T. cruzi zymograms show: a) a lack of segregation (« fixed heterozygosity » for some isozymic strains); b) a lack of recombination: each isozymic strain has its own isozymic patterns without recombination with other strains; one can observe very often the sympatric presence (even in the same Triatomine bug) of alternative alleles without recombination between them (Tibayrenc et al., 1981b and 1985). Tait (1980) inferred the existence of Mendelian sexuality for *T. brucei*.

Classification by genetic distance

Calculation of genetic distance according to Nei (1972) and Tibayrenc (1980) shows: a) the existence of 3 main groups, which correspond roughly to Miles' zymodemes (1980), but with much higher intragroup variability; b) the presence of high genetic distance values (up to 2.6), which indicates a huge level of proteic divergence (Tibayrenc and Miles, 1983; Tibayrenc and Le Ray, 1984).

Evolutive origin and biological status of *T. cruzi* isozymic strains

Isozymic data are consistent with 2 alternative hypotheses: a) Genetic distances between isozymic strains do represent an ancient divergence time, either by mitotic evolution or by real biological speciation; b) isozymic strains or at least part of them have a very recent origin by random sampling of clones within a sexuate ancestral population, and they are the genetic equivalent of individual variants without any taxonomical significance.

In the case (a), other characters of the parasite are expected to be roughly correlated to genetic distances. In the case (b), they are expected to be randomly distributed among the isozymic strains. Epidemiological data (Tibayrenc et al., 1985), pharmacological experiments (Barnabé, Tibayrenc and Dujardin, 1983) and pathogenicity studies (Brénière, in preparation), show a lack of correlation between genetic distances and other characters, and so seem to be more consistent with hypothesis (b) (Tibayrenc et al., 1985b).

Bolivian isozymic strains in domestic transmission cycles

One can observe two main isozymic strains with some lesser ones.

Heterozygous strains seem more frequent at low altitude (Tibayrenc et al., 1985a). It is easy to sample several different isozymic strains in the same house (Tibayrenc et al., 1985) and two different ones in the same Triatomine bug (Tibayrenc et al., 1985). All isozymic strains are sympatrically transmitted by the same species, *Triatoma Infestans*.

Genetic interpretation of *T. cruzi* zymograms yielded some interesting data on the biology of the parasite. Nevertheless, this theoretical approach did not yield a general theory to explain the huge biological and medical variability of Chagas' disease's causative agent.

Variabilité isoenzymatique de *T. cruzi*: portée biologique et épidémiologique.

Résumé - L'auteur passe en revue les diverses hypothéses soulevées par l'interprétation géntique de l'analyse enzymatique de *T. cruzi*.

REFERENCES

Barnabé, C., Tibayrenc, M. & Dujardin, J. P. (1983): Trypanosoma cruzi: a pharmacological comparison of some Bolivian isozymic strains. Ann. Soc. belge Méd. Trop., 63: 319-324.

Lanar, D. E., Levy, L. S. & Manning, J. E. (1981): Complexity and content of the DNA and RNA in trypanosoma cruzi. Mol. Biochem. Parasitol., 3 i 327-341.

Lemesre, J. L. & Tibayrenc, M. (1983): Trypanosoma cruzi: measurement of DNA quantity in different isoenzymic strains. Ann. Soc. belge Méd. Trop., 63: 313-317.

Miles, IV. A., Lanham, S. M., Souza, A. A. & Povoa, M. (1980): Further enzymic characters of Trypanosoma cruzi and their evaluation for strain identification. Trans. R. Soc. Trop. Med. Hyg., 74: 221-237.

Miles, M A, Toyé, P. J., Oswald, S. C. & Godfrey, D. G. (1977): The identification by isoenzyme patterns of two distinct strain-groups of Trypanosoma cruzi, circulating independently in a rural area of Brazil. Trans. R. Soc. Trop. Med. Hyg., 71 i 217-225.

Nei, M. (1972): Genetic distances between populations. Amer. Natural., 106: 283-292.

Ready, P. D. & Miles, M. A. (1980): Delimitation of Trypanosoma cruzi zymodemes by numerical taxonomy. Trans. R. Soc. Trop. Med. Hyg., 74: 238-242.

Tait, A. (1980): Evidence for diploidy and mating in trypanosomes. Nature, 287: 536-537.

Tibayrenc, M. (1980): Application of the calculations of genetic distances for flagellate systematics. Cah. ORSTOM sér. Ent. Méd. Parasitol., 18: 301-302.

Tibayrenc, M., Cariou, M. L. & Solignac, M. (1981a): Interprétation génétique des zymogrammes de flagellés des genres Trypanosoma et Leishmania. C. R. Acad. Sci. Paris, 292: 623-625.

Tibayrenc, M., Cariou, M. L., Solignac, M. & Carlier, Y. (1981b): Arguments génétiques contre l'existence d'une sexualité actuelle chez Trypanosoma cruzi implications taxonomiques. C. R. Acad. Sci. Paris, 293: 207-209.

Tibayrenc, M., Cariou, M. L., Solignac, M., Dedet, J. P., Poch, O. & Desjeux, P. (1985): New electrophoretic evidence of genetic variation in Trypanosoma cruzi. Genetica, submitted for publication.

Tibayrenc, M., Echalar, L., Brénière, F., Lemesre, J. L., Barnabé, C. & Desjeux, P. (1983): Sur le statut taxonomique et médical des souches isoenzymatiques de Trypanosoma cruzi. Considérations sur la valeur taxonomique et Immunogénique des différentes isoenzymes. C. R. Acad. Sci. Paris, 296: 721-726.

Tibayrenc M., Echalar, L., Dulardin, J. P., Poch, O. & Desjeux, P. (1984): The microdistri- 3-i cn of isoenzymic strains of Trypanosoma cruzi in Southern Bolivia. New isoenzyme profiles and further arguments against Mendelian sexuality. Trans. R. Soc. Trop. Med. Hyg., 78. 519-525.

Tibayrenc M. & Le Ray, D. (1984): General classification of the isoenzymic strains of Tripanosoma - (Schizotrypanum) cruzi and comparison with T. (S.) c. marinkellei and T. Herpetosoma) rangeli. Ann. Soc. belge Méd. trop., 64: 239-248.

Tibayrenc M & Mile, M. A. (1983): A genetic comparison between Brazilian and Bolivian zymodemes of *Trypanosoma cruzi*. Trans. B. Soc. Trop. Med. Hyg., 77: 76-83.

Tibayrenc M., Poch, O., Echalar, L., Le Pont, F., Lemesre, J. L. & Desjeux, P. (1985a): The geographical and temporal distribution of the isoenzymic strains of *Trypanosoma cruzi* in Bolivian domestic transmission cycles. Trans. D. Soc. Trop. Med. Hyg., submitted for publication.

Tibayrenc M Solignac, M., Cariou, M. L., Le Ray, D. & Desjeux, P. (1985b): Les souches izoenzymatiques de *Trypanosoma cruzi*: origine récente ou ancienne, homogène ou hétérogène? C R Acad. Sci. Paris, in press.

Toyé, P. J., Isoenzyme variation in isolates of *trypanosoma cruzi*. Trans. R. Soc. Trop. Med. Hyg., 68: 147.